

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books
 Search PubMed for [Advanced Search](#)

[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)

Display [AbstractPlus](#) Show 20 Sort By Send to

All: 1 Review: 0

1: [Ecotoxicol Environ Saf.](#) 2008 Jun; 70(2): 327-33. Epub 2008 Feb 21.

ELSEVIER Links
FULL-TEXT ARTICLE

Does Cry1Ab protein affect learning performances of the honey bee *Apis mellifera* L. (Hymenoptera, Apidae)?

[Ramirez-Romero R](#), [Desneux N](#), [Decourtye A](#), [Chaffiol A](#), [Pham-Delègue MH](#).

Instituto de Ecología A.C., Km. 2.5 Carretera Antigua a Coatepec No. 351 El Haya, 91070 Xalapa, Veracruz, Mexico.

Genetically modified Bt crops are increasingly used worldwide but side effects and especially sublethal effects on beneficial insects remain poorly studied. Honey bees are beneficial insects for natural and cultivated ecosystems through pollination. The goal of the present study was to assess potential effects of two concentrations of Cry1Ab protein (3 and 5000 ppb) on young adult honey bees. Following a complementary bioassay, our experiments evaluated effects of the Cry1Ab on three major life traits of young adult honey bees: (a) survival of honey bees during sub-chronic exposure to Cry1Ab, (b) feeding behaviour, and (c) learning performance at the time that honey bees become foragers. The latter effect was tested using the proboscis extension reflex (PER) procedure. The same effects were also tested using a chemical pesticide, imidacloprid, as positive reference. The tested concentrations of Cry1Ab protein did not cause lethal effects on honey bees. However, honey bee feeding behaviour was affected when exposed to the highest concentration of Cry1Ab protein, with honey bees taking longer to imbibe the contaminated syrup. Moreover, honey bees exposed to 5000 ppb of Cry1Ab had disturbed learning performances. Honey bees continued to respond to a conditioned odour even in the absence of a food reward. Our results show that transgenic crops expressing Cry1Ab protein at 5000 ppb may affect food consumption or learning processes and thereby may impact honey bee foraging efficiency. The implications of these results are discussed in terms of risks of transgenic Bt crops for honey bees.

PMID: 18206234 [PubMed - indexed for MEDLINE]

Display [AbstractPlus](#) Show 20 Sort By Send to

Related Articles

Effects of transgenic corn and Cry1Ab protein on the nematode, *Caenorhabditis elegans*. [Ecotoxicol Environ Saf. 2008]

Learning performances of honeybees (*Apis mellifera* L) are differentially affected by imidacloprid according to the season. [Pest Manag Sci. 2003]

A meta-analysis of effects of Bt crops on honey bees (Hymenoptera: Apidae). [PLoS ONE. 2008]

Review Behavioral studies of learning in the Africanized honey bee (*Apis mellifera* L.). [Brain Behav Evol. 2002]

Review Climate change: impact on honey bee populations and diseases. [Rev Sci Tech. 2008]

» See Reviews... | » See All..

Recent Activity

[Turn Off](#) [Clear](#)

Does Cry1Ab protein affect learning performances of the honey bee *Apis mellifera* L. (Hymen...

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)