



Pentagon Reports: Fast. Definitive. Complete.

[Home](#) [About Us](#) [Contact Us](#) [View Cart](#) [My Account](#) [FAQ](#)

username

**LOGIN**

[New Account »](#)  
[Forgot Password?](#)

Aluminum oxide



[Advanced Search »](#)

Ads by Google

[Aluminum Powder](#)

German  
Blackhead \$17/LB  
Atomized \$12,  
Indian Blackhead  
\$17

[www.HighQualityChems.com](http://www.HighQualityChems.com)

[US Army - Official Site](#)

Earn \$2,000 when  
you refer someone  
to the Army.

Details inside!

[www.army.mil](http://www.army.mil)

[Saturn Official Site](#)

Come See the Full  
Lineup of Saturn  
Vehicles with  
Great Gas  
Mileage.

[www.Saturn.com/GasMile](http://www.Saturn.com/GasMile)

[Infrared Spectroscopy](#)

Search Thousands  
of Catalogs for  
Infrared

Spectroscopy

[www.globalspec.com](http://www.globalspec.com)



Chemistry ▾ [Inorganic Chemistry](#)

**Laboratory Studies of Al(2)O(3)-NO(x) Aerosols**

Authors: [Robert Disselkamp](#); [John R. Edwards](#); [Daniel Pilson](#); [Tyrrel W. Smith Jr.](#); TRW SPACE AND ELECTRONICS GROUP  
[REDONDOBEACH CA](#)

**Abstract:** Laboratory experiments were performed to investigate the chemistry of **aluminum oxide** (gamma Al(2)O(3)) aerosol samples upon exposure to NO(x) (NO(x) is NO, NO(2), etc..) gases. Static aerosol samples were generated in an aerosol chamber and studied at temperatures ranging from 298 to 183 K. Fourier-transform infrared (FTIR) absorption spectroscopy was used to study the aerosol samples over time. Each aerosol was created using the same procedure. First, a reactant gas species, NO or NO(2), was added to the chamber and infrared spectra were collected over a 20-minute time interval to characterize heterogeneous reactions occurring on the chamber walls. Next, an **aluminum oxide** aerosol was generated by expanding powder into the chamber using nitrogen gas at high pressure. Infrared spectra were then collected at 6-minute intervals for at least 100 minutes to characterize Al(2)O(3)-NO(x) chemistry. The experiments reported on in this paper enabled a quantitative characterization of both the rate of reactant gas uptake and product formation processes to be performed. A quantitative (i.e., stoichiometric) analysis of reactant gas depletion and product gas formation enabled elementary reactions involving **aluminum oxide** surface hydroxyl sites and NO(x) species to be proposed. (2 tables, 2 figures, 6 refs.)

Adobe PDF - \$8.95

Printed Format - \$23.95



Please check the box for the format you wish to order.

[Shipping Terms](#)  
[About Electronic Delivery](#)

[Email This Abstract](#)

**Limitations:**  APPROVED FOR PUBLIC RELEASE

**Description:** Technical rept.

**Pages:** 13

**Report Date:** 30 SEP 1999

**Contract Number:** F09603-95-D-0176-0007

**Report Number:** A504414



**Keywords relating to this report:**

- ✦ [ABSORPTION SPECTRA](#)
- ✦ [AEROSOLS](#)
- ✦ [ALUMINUM OXIDES](#)
- ✦ [CHAMBERS](#)
- ✦ [CHEMICAL REACTIONS](#)
- ✦ [DEPLETION](#)
- ✦ [FOURIER SPECTROSCOPY](#)
- ✦ [HIGH PRESSURE](#)
- ✦ [INFRARED SPECTRA](#)
- ✦ [INFRARED SPECTROSCOPY](#)
- ✦ [LABORATORY TESTS](#)
- ✦ [NITROGEN OXIDES](#)
- ✦ [REACTANTS\(CHEMISTRY\)](#)
- ✦ [REACTIVE GASES](#)
- ✦ [STOICHIOMETRY](#)

[« Back to search](#)

[Home](#) | [About Us](#) | [Contact Us](#) | [View Cart](#) | [Customer Service](#) | [Shipping Terms](#) | [Advanced Search](#) | [Privacy Policy](#) | [Restrictions on PDF Usage](#)

© 2001-2008 Storming Media LLC. All rights reserved.