

username

LOGIN

[New Account »](#)
[Forgot Password?](#)

CHAFF

GO

[Advanced Search »](#)

[Oceanography and Atmospheric Sci](#) » [Atmospheric Physics](#)

WIND MEASUREMENTS IN THE SUBPOLAR MESOPAUSE REGION

Authors: [James E. Morris](#); [ARMY ELECTRONICS COMMAND WHITE SANDS MISSILE RANGE NM ATMOSPHERIC SCIENCES LAB](#)

Abstract: Mesospheric wind data obtained with a new high altitude Loki system during the summer of 1966 over Fort Greely, Alaska, are presented. Soundings, utilizing very light **chaff** as a wind sensor, were scheduled near noon and midnight for a 50-hour period. These data are from a sparsely sampled region of the atmosphere. The diurnal variations and the high velocities observed give valuable information regarding noctilucent clouds, atmospheric tidal oscillations, and the mean summer flow near the subpolar mesopause.

Adobe PDF - \$18.95

Printed Format - \$20.95

ADD TO CART

Please check the box for the format you wish to order.

Limitations: APPROVED FOR PUBLIC RELEASE

Pages: 30

Report Date: JAN 1967

Report Number: 0318056

[Shipping Terms](#)
[About Electronic Delivery](#)

[Email This Abstract](#)

Keywords relating to this report:

- ✦ [*Mesosphere](#)
- ✦ [*RADAR CONFUSION REFLECTORS](#)
- ✦ [ALASKA](#)
- ✦ [ATMOSPHERES](#)
- ✦ [DETECTORS](#)
- ✦ [HIGH ALTITUDE](#)
- ✦ [NOCTILUCENT CLOUDS](#)
- ✦ [PERIODIC VARIATIONS](#)
- ✦ [SAMPLING](#)
- ✦ [VELOCITY](#)
- ✦ [WIND](#)

[« Back to search](#)