

News

Articles

Videos

Images

Books

[Health & Medicine](#)
[Mind & Brain](#)
[Plants & Animals](#)
[Earth & Climate](#)
[Space & Time](#)
[Matter & Energy](#)
[Computers & Math](#)
[Fossils & Ruins](#)

[Share](#)
[Blog](#)
[Cite](#)
[Print](#)
[Email](#)
[Bookmark](#)

Science News

Oceanography: Low Summer Iron Availability Limits Biological Production In The High-Latitude North Atlantic

ScienceDaily (July 26, 2009) — Southampton scientists have demonstrated an unexpected role of iron in regulating biological production in the high-latitude North Atlantic. Their findings have important implications for our understanding of ocean-climate interactions.

See Also:

Earth & Climate

- [Global Warming](#)
- [Geography](#)
- [Oceanography](#)
- [Climate](#)
- [Environmental Issues](#)
- [Earth Science](#)

Reference

- [Phytoplankton](#)
- [Plankton](#)
- [Carbon cycle](#)
- [Ocean acidification](#)

Tiny plant-like organisms called phytoplankton dominate biological production in the sunlit surface waters of the world's oceans and, through the process of photosynthesis, sequester large amounts of atmospheric carbon dioxide. A proportion of the carbon is exported to the deep ocean, and because carbon dioxide is a greenhouse gas, this so-called 'biological carbon pump' helps prevent runaway global climate warming.

Iron is an essential micro nutrient for phytoplankton growth. In high-nutrient, low-chlorophyll (HNLC) oceanic regions, phytoplankton growth is limited by low iron

availability. Classical HNLC regions, which account for about a third of the world's oceans, include the Southern Ocean and the subpolar North Pacific.


In contrast, it has been widely assumed that iron supply does not limit biological production of the high-latitude (>50 degrees N) North Atlantic Ocean. Here, winter cooling causes the sinking of nutrient-depleted surface waters and their replacement by deep nutrient-rich water. This winter 'overturning' replenishes surface water nutrients, and in the spring, when light intensities increase, a large phytoplankton bloom develops, leading to high rates of carbon export.

However, in many regions of the open North Atlantic, including the Iceland and Irminger Basins, residual amounts of nitrate persist into the summer period, after the spring bloom has ceased. This represents an inefficiency of the biological carbon pump that is potentially of global significance to the partitioning of carbon between the atmosphere and ocean.

Phytoplankton are grazed upon and some of the larger phytoplankton species such as diatoms with low grazing mortality are susceptible to silicate shortage. Traditionally, it has been believed that these factors, acting in concert, might be sufficient to terminate the spring bloom leaving some nitrate unused. However, there have been indications that phytoplankton might simply run out of iron before they are able to exploit any remaining other nutrients.

Now a team of scientists from the National Oceanography Centre, Southampton, have tested this 'iron limitation hypothesis' for the high-latitude North Atlantic Ocean. Their measurements were performed on a cruise aboard the RRS Discovery within the central Iceland Basin during the summer of 2007, and the findings are published this month in the scientific journal *Global Biogeochemical Cycles*.

The researchers found that the concentration of dissolved iron in surface waters was very low, as was biological production, despite the presence of residual nitrate. Experimental addition of iron to bottles containing seawater samples increased photosynthetic efficiency, chlorophyll concentrations, and growth of several types of phytoplankton, including the ubiquitous *Emiliania huxleyi* a coccolithophore



Emiliania huxleyi, a type of phytoplankton. (Credit: Image courtesy of National Oceanography Centre, Southampton)

Ads by Google

Climate change

UPS Strives to Protect the Earth for our People and our Community.
www.sustainability.ups.com

Global Warming = MONEY

Climate Change who Profits?
What are the facts?
NexPlanRecycling.com

Objet 3D Printing Systems

Create & Print 3D Prototypes.
Watch our Demo and be Amazed!
Objet.com

Free Trial- Phytoplankton

Highest Quality Pure Multi-Strain
Phytoplankton (1 Bottle \$29.95)
www.MarinePhytoplankton5000.com

Related Stories



Iron Controls Patterns Of Nitrogen Fixation In The Atlantic

(Nov. 2, 2009) — Scientists have discovered that interactions between iron supply, transported through the atmosphere from deserts, and large-scale oceanic circulation control the availability of a crucial nutrient, ... [> read more](#)



Oceanic Seesaw Links Northern And Southern Hemisphere During Abrupt Climate Change During Last Ice Age

(Mar. 3, 2009) — Very large and abrupt changes in temperature recorded over Greenland and across the North Atlantic during the last Ice Age were actually global in extent, according to new research. The research ... [> read more](#)



Dust Storms In Sahara Desert Sustain Life In Atlantic Ocean

(July 19, 2008) — Saharan dust storms help sustain life over extensive regions of the North Atlantic Ocean. Scientists mapped the distribution of nutrients including phosphorous and nitrogen and investigated how ... [> read more](#)

[Understanding Sea Temperature Atmospheric](#)

Just In:

[First Data from the Large Hadron Collider](#)

Science Video News



Global Warming Equals Stronger Hurricanes

Climate change experts studying hurricanes documented a 35-year warming trend in ocean surface temperature and linked it to larger hurricanes. The ... [> full story](#)

[Meteorologists Discover How Monsoon Affects Summer Weather](#)

[Improving Accuracy Of Hurricane Forecasts](#)

[Statisticians Show that High Altitude Makes Hits Longer](#)

[more science videos](#)



Breaking News

... from [NewsDaily.com](#)

[NASA hauls shuttle to launch pad, despite cold](#)

[Study turns up 10 autism clusters in California](#)

[Tombs to lift lid on Egypt's ancient middle class](#)

[One in 10 births around world premature: WHO](#)

[Relic of Antarctica's first plane found on ice-edge](#)

[more science news](#)

In Other News ...

[A goodbye kiss triggered New Jersey airport scare](#)

[Pelosi says Congress close to health deal](#)

["Shock" U.S. report due on attempted airline attack](#)

[Three dead in Missouri factory shooting](#)

biological production in the high-latitude North Atlantic. This has only previously been suspected, but helps explain why the spring phytoplankton bloom does not continue well into the summer and why residual amounts of nitrate remain unused."

The central Iceland Basin receives little iron input from continental sources or from atmospheric dust, and some iron is also lost through detrital sinking. Moreover, the new findings suggest that iron brought to the surface during winter overturning is insufficient to support maintenance of the phytoplankton bloom into the summer.

Nielsdottir, a research student at the University of Southampton's School of Ocean and Earth Science based at the National Oceanography Centre, said: "In effect, the high-latitude North Atlantic is a seasonal HNLC region, whereas classic, HNLC regions such as the Southern Ocean remain in this condition throughout the year."

The failure of the phytoplankton community to exploit residual nitrate remaining in the summer reduces the effectiveness of the biological carbon pump. "This is important," says Nielsdottir, "because the high-latitude North Atlantic is second only to the Southern Ocean in its potential to lower atmospheric carbon dioxide and un used nitrate in the surface highlight the potential for even higher CO₂ drawdown, high levels of which are an important cause of global climate warming."

The work was supported by the National Oceanography Centre with a PhD to MCN, Natural Environment Research Council, Oceans2025 and the Faroese Ministry of Interior and Law. The authors are Maria Nielsdottir, C. Mark Moore, Richard Sanders, Daria Hinz and Eric Achterberg, University of Southampton's School of Ocean and Earth Science based at the National Oceanography Centre, Southampton.

Email or share this story: | [More](#)

Story Source:

Adapted from materials provided by [National Oceanography Centre, Southampton \(UK\)](#).

Journal Reference:

Nielsdóttir, M. C., C. M. Moore, R. Sanders, D. J. Hinz, and E. P. Achterberg. **Iron limitation of the postbloom phytoplankton communities in the Iceland Basin.** *Biogeochem. Cycles*, 23, GB3001 DOI: [10.1029/2008GB003410](#)

Need to cite this story in your essay, paper, or report? Use one of the following formats:

- ☒ APA National Oceanography Centre, Southampton (UK) (2009, July 26). Oceanography: Low Summer Iron Availability Limits Biological Production In The High-latitude North Atlantic. *ScienceDaily*. Retrieved January 8, 2010, from <http://www.sciencedaily.com/releases/2009/07/090707111711.htm>
- ☐ MLA

Note: If no author is given, the source is cited instead.

studies have found that atmospheric anomalies associated with a ... > [read more](#)



Study Provides First-Ever Look At Combined Causes Of North Atlantic And Arctic Ocean Freshening

(Aug. 25, 2006) — A new analysis of 50 years of changes in freshwater inputs to the Arctic Ocean and North Atlantic may help shed light on what's behind the recently observed freshening of the North Atlantic Ocean. ... > [read more](#)



Understanding Ocean Climate

(Dec. 13, 2009) — High-resolution computer simulations are helping to describe the inflow of North Atlantic water to the Arctic Ocean and how this influences ocean ... > [read more](#)



Mapping Nutrient Distributions Over The Atlantic Ocean

(Dec. 30, 2009) — Large-scale distributions of two important nutrient pools -- dissolved organic nitrogen and dissolved organic phosphorus have been systematically mapped for the first time over the Atlantic Ocean in ... > [read more](#)

Climate Change Leads To Major Decrease In Carbon Dioxide Storage (Apr. 10, 2009) — The North Atlantic Ocean is one of the Earth's tools to offset natural carbon dioxide emissions. In fact, the 'carbon sink' in the North Atlantic is the primary gate for carbon dioxide entering the ... > [read more](#)

Ads by Google

The Climate Challenge

101 Solutions to Global Warming
Act now to help Stop Climate Change
[www.newsociety.com](#)

Cryogenic Specimen Moving

Experienced lab relocators with climate-controlled moving vans.
[www.AcceleratedMoving.com](#)

Research climate change

Learn whats new without the clutter
Free service with daily email alert
[topikality.com](#)

Pure Marine Phytoplankton

RePhytoLize & Oceans Alive
100% Pure, Value - Free Shipping!
[PureHealingFoods.com](#)

Search ScienceDaily

Number of stories in archives: 79,715

Blackwater case

Iran to try five people over December unrest

Nigeria says plane suspect was properly screened

[more top news](#)

Copyright Reuters 2008. See [Restrictions](#).

Free Subscriptions

... from ScienceDaily

Get the latest science news with our free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

[Email Newsletters](#)

[RSS Newsfeeds](#)

Feedback

... we want to hear from you!

Tell us what you think of the new ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

Your Name:

Your Email:

Comments:

Click button to submit feedback:

[Send It](#)

Find with keyword(s):

[Search](#)

Enter a keyword or phrase to search ScienceDaily's archives for related news topics, the latest news stories, reference articles, science videos, images, and books.