



## Institutional Login

## Welcome!

To use the personalized features of this site, please [log in](#) or [register](#).

If you have forgotten your username or password, we can [help](#).

## My Menu

[Marked Items](#)

[Alerts](#)

[Order History](#)

## Saved Items


[All](#)

[Favorites](#)

Content Types Subject Collections

English

## Book Chapter



large version

## Wheat Breeding for Soil Acidity and Aluminum Toxicity

Book Series Developments in Plant Breeding  
 ISSN 1381-673X  
 Volume Volume 12  
 Book Wheat Production in Stressed Environments  
 Publisher Springer Netherlands  
 DOI 10.1007/1-4020-5497-1  
 Copyright 2007  
 ISBN 978-1-4020-5496-9 (Print) 978-1-4020-5497-6 (Online)  
 DOI 10.1007/1-4020-5497-1\_48  
 Pages 387-394  
 Subject Collection Biomedical and Life Sciences  
 SpringerLink Date Wednesday, May 16, 2007

**Add to marked items**  
 Add to shopping cart  
 Add to saved items  
 Recommend this chapter

PDF (240.6 KB) [Free Preview](#)

Developments in Plant Breeding

Wheat Production in Stressed Environments  
 Proceedings of the 7th International Wheat Conference, 27  
 November–2 December 2005, Mar del Plata, Argentina

10.1007/1-4020-5497-1\_48

H. T. Buck, J. E. Nisi and N. Salomón

### E. von Baer<sup>4</sup>

(4) Campex – Semillas Baer and AVELUP Fundo El Hualle s/n, Cajón-Temuco, Chile

#### Abstract

The soils of southern and central-southern Chile where the national wheat production is concentrated are of volcanic origin, with low pH and with high exchangeable aluminum. Soils contain large amounts of phosphorous in organic form, however available phosphorous is low. The high content of free Al negatively affects the root growth of wheat varieties depending on their Al-sensitivity or Al-tolerance. The mechanism of tolerance is exclusion of toxic Al as a consequence of root excretion of chelant organic acids; roots of tolerant varieties are also protected by higher root mycorrhizal colonization. Crosses between sensitive and tolerant material and subsequent selection under low pH and high Al have produced high yielding tolerant cultivars. This paper outlines the development of tolerant wheat varieties, describes the germplasm used and discusses the selection process including the use of growth chambers, greenhouse assays and field experiments

**Keywords** Al-tolerant varieties - citric acid - rhizosphere - mycorrhizal colonization - segregant material

**E. von Baer**  
 Email: [infoseedsbaer@semillasbaer.cl](mailto:infoseedsbaer@semillasbaer.cl)

[Fulltext Preview \(Small, Large\)](#)

Find [more options](#)

- Within all content
- Within this book series
- Within this book

#### Export this chapter

Export this chapter as RIS | Text

Ads by Google

#### [Aluminum PegBoard X2™](#)

World Leader in  
 Aluminum PegBoard,  
 Strong, Sleek, Load  
 Rated, \$6.99/sf  
[www.diamondlifegear.com/pegbox](http://www.diamondlifegear.com/pegbox)

#### [Aluminum Castings](#)

Intricately Cored  
 Aluminum Castings US  
 Aluminum Sand Casting  
 Foundry  
[www.buddybarcasting.com](http://www.buddybarcasting.com)

#### [Oral Chelation Works](#)

Remove Aluminum & 15  
 Heavy Metals Safely &  
 Effectively From the Body  
[www.scienceformulas.com](http://www.scienceformulas.com)

#### [Aluminum Custom Extrusion](#)

Custom Aluminum  
 Extrusion Shapes &  
 Design-Assistance and  
 Fabrication  
[www.AmericanDouglasMetals.com](http://www.AmericanDouglasMetals.com)

#### [Machined Aluminum Parts](#)

3D CAD to CNC  
 machined part as fast as  
 1 business day. Online  
 quotes  
[www.firstcut.com](http://www.firstcut.com)

## WHEAT BREEDING FOR SOIL ACIDITY AND ALUMINUM TOXICITY

E. VON BAER

*Campex – Semillas Baer and AVELUP, Fundo El Hualle s/n, Cajón-Temuco, Chile*  
*E-mail: info@semillasbaer.cl*

**Abstract:** The soils of southern and central-southern Chile where the national wheat production is concentrated are of volcanic origin, with low pH and with high exchangeable aluminum. Soils contain large amounts of phosphorous in organic form, however available phosphorous is low. The high content of free Al negatively affects the root growth of wheat varieties depending on their Al-sensitivity or Al-tolerance. The mechanism of tolerance is exclusion of toxic Al as a consequence of root excretion of chelant organic acids; roots of tolerant varieties are also protected by higher root mycorrhizal colonization. Crosses between sensitive and tolerant material and subsequent selection under low pH and high Al have produced high yielding tolerant cultivars. This paper outlines the development of tolerant wheat varieties, describes the germplasm used and discusses the selection process including the use of growth chambers, greenhouse assays and field experiments

**Keywords:** Al-tolerant varieties, citric acid, rhizosphere, mycorrhizal colonization, segregant material

### INTRODUCTION

The soils of southern and central-southern Chile (35°S to 42°S), are of volcanic origin and characterized by high acidity (pH range from 4.5 to 6) and high aluminum saturation approaching 30% without lime and 0–18% with applied lime. Nutrient availability including P is low. These constrictive characteristics are known as the soil acidity complex which often creates a chemical barrier reducing root growth and development; these roots become inefficient in absorbing nutrients and water (Reynolds et al. 2001).

Despite this, volcanic soils contain high levels of total P (2,000 to 4,000 mg kg<sup>-1</sup>) but the availability of this nutrient is low (5–20 ppm). The unavailability of P to

387

*H.T. Buck et al. (eds.), Wheat Production in Stressed Environments, 387–393.*  
© 2007 Springer.

References secured to subscribers.

© Springer. Part of Springer Science+Business Media  
Privacy, Disclaimer, Terms and Conditions, © Copyright Information

[Privacy Policy](#)

Remote Address: 66.81.65.158 • Server: mpweb16  
HTTP User Agent: Mozilla/4.0 (compatible; MSIE 7.0; Windows NT 6.0; SLCC1; .NET CLR 2.0.50727; Media Center PC 5.0; .NET CLR 3.0.04506)