

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books  
 Search PubMed for    [Advanced Search](#)

[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)

Display **AbstractPlus** Show **20** Sort By Send to

1: [Ecotoxicol Environ Saf.](#) 2008 Jun; 70(2): 334-40. Epub 2008 Feb 20.

**ELSEVIER** Links  
 FULL-TEXT ARTICLE

**Effects of transgenic corn and Cry1Ab protein on the nematode, *Caenorhabditis elegans*.**

[Höss S](#), [Arndt M](#), [Baumgarte S](#), [Tebbe CC](#), [Nguyen HT](#), [Jehle JA](#).

Ecossa (Ecological Sediment and Soil Assessment), Giselastr. 6, 82319 Starnberg, Germany. hoess@ecossa.de

The effects of the insecticidal Cry1Ab protein from *Bacillus thuringiensis* (Bt) on the nematode, *Caenorhabditis elegans*, were studied with soil from experimental fields cultivated with transgenic Bt corn (MON810) and with trypsinized Cry1Ab protein expressed in *Escherichia coli*. The content of Cry1Ab protein was above the detection limit of an ELISA test in only half of the soil samples obtained from transgenic plots, ranging from 0.19 to 1.31 ng g(-1) dry weight. In a laboratory bioassay, *C. elegans* was exposed to rhizosphere and bulk soil from fields with isogenic or transgenic corn or to solutions of Cry1Ab protein (0, 24, 41, 63, 118, and 200 mg l(-1)) over a period of 96 h, with growth and reproduction serving as the test parameters. Nematode reproduction and growth were significantly reduced in rhizosphere and bulk soil of Bt corn compared with soil from isogenic corn and were significantly correlated with concentrations of the Cry1Ab protein in the soil samples. Moreover, the toxicity of pure Cry1Ab protein to the reproduction and growth of *C. elegans* was concentration-dependent. As significant inhibition occurred at relatively high concentrations of the Cry1Ab protein (41 mg l(-1)), the effects of the soil samples from Bt corn could not be assigned directly to the toxicity of the Cry1Ab protein. The results demonstrate that bioassays with the nematode, *C. elegans*, provide a promising tool for monitoring the potential effects of Bt toxins in aqueous medium and soils.

PMID: 18068780 [PubMed - indexed for MEDLINE]

Display **AbstractPlus** Show **20** Sort By Send to

**Related Articles**

Field studies on the environmental fate of the Cry1Ab Bt-toxin produced by transgenic maize (MON810) and its effect on bacterial communities in the maize rhizosphere. *Microb Ecol.* 2005

Occurrence and persistence of *Bacillus thuringiensis* (Bt) and transgenic Bt corn cry1Ab gene from an aquatic environment. *[Ecotoxicol Environ Saf. 2007]*

DiPel-selected *Ostrinia nubilalis* larvae are not resistant to transgenic corn expressing *Bacillus thuringiensis* Cry1Ab. *PLoS One.* 2007

*Review* Monitoring and adaptive resistance management in Australia for Bt-cotton: current status and future challenges. *PLoS One.* 2007

*Review* [Potential effects of Bt transgenic rice on soil micro-ecosystem] *[Ying Yong Sheng Tai Xue Bao. 2005]*

» See Reviews... | » See All...

**Recent Activity**

Effects of transgenic corn and Cry1Ab protein on the nematode, *Caenorhabditis elegans*.

Does Cry1Ab protein affect learning performances of the honey bee *Apis mellifera* L. (Hymen...

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)